

SUBSCRIBE

www.naturefresh.co.za

Fresh Talk

FRESH Talk

Natural Solutions to
Health Questions

NATURE
FRESH



EPISODE 23:

SUBSCRIBE

Fresh Talk

NATURE
FRESH

www.naturefresh.co.za

Can Health Foods be FAKE?

*Secrets of S.I.B.O and Why eating fibre is
destroying your gut...*

SYNTHESIS OF GLUTEN PATHOLOGY, COMMERCIAL GLUTEN-FREE PITFALLS, AND DIETARY PROTOCOLS

Executive Summary : Research by Dr Eric Berg and Sue Visser

Gluten—a protein component found in wheat, rye, barley, and related grains—presents a complex clinical picture affecting gastrointestinal, neurological, endocrine, and systemic health. Contrary to popular perception, human physiology lacks the enzymes required to fully digest gluten protein, triggering gut inflammation and systemic health complications across diverse populations. The modern addition of refined gluten to processed foods has exacerbated chronic conditions, linking gluten exposure to over 55 distinct diseases, ranging from celiac disease and leaky gut syndrome to severe psychiatric disorders and autoimmune conditions via molecular mimicry.

A critical finding synthesized across the source materials is the commercial fallacy of processed “gluten-free” (GF) substitute foods. Manufactured GF alternatives frequently replace wheat with high-glycemic refined starches (e.g., tapioca, corn, potato, and rice starches), seed oils, and added sugars. These substitutes lack adequate protein, spike blood sugar and insulin levels more dramatically than table sugar, and foster gut dysbiosis, including Small Intestinal Bacterial Overgrowth (SIBO). Furthermore, ingredients like rice flour introduce secondary risks, such as arsenic bioaccumulation.

Optimal health management necessitates a fundamental distinction between merely avoiding gluten and adopting a holistically grain-free, nutrient-dense diet tailored to individual bio-individuality, such as blood type secretor status.

OFFICIAL DISCLAIMER: All material provided is only information and may not be construed as medical advice. Only a doctor or an appropriate health practitioner is qualified to make a diagnosis and provide the necessary treatment and medications. Products discussed are not intended to diagnose, cure or prevent any form of disease.

SUBSCRIBE

www.naturefresh.co.za

Fresh Talk

FRESH Talk

Natural Solutions to Health Questions

with NATURE FRESH

Correcting gluten-induced damage requires identifying hidden sources of exposure, eliminating refined starches and grains, addressing gut dysbiosis with targeted Whole Foods or temporary elimination diets (e.g., carnivore protocol), and re-establishing beneficial microbiome populations using fermented foods.

The Dual Pathology of Wheat, Gluten, and Wheat Germ Agglutinin (WGA)

Digestibility and Structural Toxicity

Gluten is the primary protein component of wheat and related grains. It represents the single protein structure that the human digestive tract cannot fully break down. Modern agricultural and food-processing practices have drastically elevated gluten concentrations in standard diets, driving chronic inflammation across the gastrointestinal (GI) tract and systemic organs.

Beyond gluten itself, whole wheat—including sprouted and organic varieties—contains high concentrations of Wheat Germ Agglutinin (WGA), a glycoprotein classified as a lectin. WGA possesses a unique molecular architecture reinforced by disulfide bonds—the same chemical bonds that confer resilience to vulcanized rubber and human hair. Consequently, WGA resists thermal and enzymatic breakdown across extreme pH ranges, leading to bioaccumulation within human tissues.

COMPONENTS OF WHEAT TOXICITY	
Component	Pathological Mechanism / Impact
Gluten Protein	Indigestible protein structure; damages intestinal villi, increases gut permeability, induces celiac.
Wheat Germ Agglutinin (WGA)	Resilient disulfide-bonded lectin; bioaccumulates, breaches blood-brain barrier, disrupts endocrine axis.
Gliadin	Alcohol-soluble protein fraction triggering immune sensitivity, celiac antibodies, and “addictive” response
Gliadorphin (Exorphins)	Opioid peptides formed during gluten digestion; mask internal pain, driving allergenic food addiction.
Excitotoxic Amino Acids	High glutamic and aspartic acid content leading to nerve cell receptor over-activation and brain injury.

OFFICIAL DISCLAIMER: All material provided is only information and may not be construed as medical advice. Only a doctor or an appropriate health practitioner is qualified to make a diagnosis and provide the necessary treatment and medications. Products discussed are not intended to diagnose, cure or prevent any form of disease.



Pathological Mechanisms of WGA and Gluten

WGA and gluten fractions exert systemic toxicity across multiple physiological domains:

- * Pro-inflammatory & Intestinal Epithelial Damage: WGA stimulates the synthesis of pro-inflammatory cytokines within intestinal and immune cells, promoting chronic thin-gut inflammation. It causes shedding of the brush border membrane, reduces mucosal surface area, accelerates enterocyte death, and depletes heat shock protective proteins.
- * Autoimmunity via Molecular Mimicry: The structural geometry of gluten proteins closely resembles human tissues, such as the thyroid gland and the myelin sheath. Immune antibodies generated against gluten erroneously attack target host tissues, manifesting as Hashimoto's thyroiditis, Multiple Sclerosis (MS), or rheumatoid complications.
- * Immunotoxicity: WGA induces thymus atrophy in animal models, while anti-WGA antibodies cross-react with human tissue proteins, distinct from standard gluten-induced celiac autoimmunity.
- * Neurotoxicity & Brain Barrier Disruption: WGA crosses the blood-brain barrier via adsorptive endocytosis, pulling co-transported toxic substances into the central nervous system. It binds to the myelin sheath and inhibits Nerve Growth Factor (NGF), impairing neuronal survival and regeneration.
- * Cardiotoxicity: WGA impairs platelet endothelial cell adhesion molecule-1, obstructing normal tissue regeneration and the safe clearance of neutrophils from blood vessels.
- * Endocrine Disruption: WGA binds leptin receptors within the hypothalamus, driving leptin resistance, uninhibited appetite, and weight gain. It further interferes with pancreatic secretin release, causing digestive insufficiency and pancreatic hypertrophy.
- * Excitotoxicity: Wheat contains elevated levels of glutamic acid and aspartic acid. Excess exposure over-activates N-methyl-D-aspartate (NMDA) nerve cell receptors, inducing calcium-mediated neuronal injury associated with MS, Alzheimer's disease, Huntington's disease, epilepsy, ADHD, and migraines.

Systemic Health Manifestations

Unmitigated gluten and wheat consumption contributes to widespread symptomatic conditions:

- * Joint and Skeletal Deterioration: Gliadin and wheat lectins rob joint tissue of native glucosamine, precipitating stiffness and joint degradation. Furthermore, wheat bran contains indigestible phytic acid, which binds systemic calcium. The body leaches skeletal calcium to neutralize the resulting systemic acidity, predisposing individuals to osteoporosis and acidic urine-linked kidney stones.
- * Urological Dysfunction: Ingestion of wheat and gluten frequently triggers involuntary bladder wall spasms, drastically reducing bladder holding capacity. This manifests as chronic frequent urination, nocturia, or pediatric bedwetting.
- * Refractory Anaemia: Chronic gluten consumption flattens microscopic intestinal villi, impairing nutrient absorption. Individuals with undiagnosed gluten sensitivity routinely present with severe iron-deficiency anaemia that remains completely non-responsive to oral iron supplementation until all gluten sources are removed.
- * Impaired Vitamin Assimilation: Gluten intolerance severely inhibits the intestinal assimilation of essential fat-soluble nutrients, notably Vitamin D, even in individuals receiving ample sunlight exposure.

The Fallacy of Commercial "Gluten-Free" Products

The Nutritional Reality of GF Processing

Commercial marketing often equates "gluten-free" labeling with superior health quality. In reality, mass-market GF substitute products are frequently more destructive to metabolic and gastrointestinal health than standard grain products.

OFFICIAL DISCLAIMER: All material provided is only information and may not be construed as medical advice. Only a doctor or an appropriate health practitioner is qualified to make a diagnosis and provide the necessary treatment and medications. Products discussed are not intended to diagnose, cure or prevent any form of disease.

To replicate the structural elasticity of gluten, manufacturers employ industrial extraction techniques on grains, utilizing chemical solvents, synthetic enzymes, and mineral acids to strip away protein fractions. The resulting isolate is a highly processed, nutrient-depleted base.

COMMERCIAL GLUTEN-FREE INGREDIENTS VS. HEALTH RISKS	
Commercial Substitute	Biological / Pathological Consequence
Refined Starches (Corn, Potato, Tapioca)	High glycemic index; spikes blood glucose and insulin higher than table sugar; accelerates fatty liver.
Cornstarch	Highly refined starch (used industrially for clothing stiffening); fuels pathogenic bacterial overgrowth.
Rice Flour	High carbohydrate payload; hyper-accumulates environmental heavy metal arsenic from soil.
Bioengineered Ingredients Industrial Seed Oils	Exposure to Genetically Modified Organisms (GMOs) and pro-inflammatory processed fats.
Commercial Binders (Xanthan / Guar Gum)	Alters nutrient absorption; causes intestinal cramps, fermentation, severe bloating, and gas in sensitive items.

Metabolic and Blood Sugar Impact

Starches are long chains of glucose molecules. On the glycemic index, refined starches (such as potato starch, cornstarch, food starch, and tapioca starch) rank higher than refined table sugar, causing sharper blood glucose surges and exaggerated insulin spikes. This chronic hyper-insulinaemia promotes systemic inflammation, rapid weight gain, visceral adiposity, and cellular insulin resistance.


Additionally, commercial GF bakery goods (e.g., cookies, breads, crackers, granola bars) are notoriously low in protein—the primary macronutrient responsible for triggering satiety signals. Consequently, consumption of low-protein, highly refined GF starches fails to satisfy hunger, leading to persistent food cravings, overeating, and metabolic instability.

Microbiome Disruption and SIBO

Pathogenic microflora thrive on refined simple sugars and starches, whereas beneficial commensal gut bacteria depend on indigestible prebiotic plant fibers. Dumping high-glycemic GF carbohydrates into the GI tract disrupts microbial ecology, resulting in dysbiosis:



1. Toxic Byproduct Cascade: As overfed pathogenic bacteria die off, they release toxic metabolic endotoxins. This triggers localized intestinal inflammation that radiates through the portal system, causing hepatic inflammation, non-alcoholic fatty liver disease, and systemic dermatological manifestations (e.g., severe dermatitis, eczema, acne).
2. Small Intestinal Bacterial Overgrowth (SIBO): Pathogenic organisms migrate upward and proliferate inside the small intestine, competing with the host for vital vitamins and minerals.
3. The Fiber Paradox: When SIBO is present, consuming high-fiber foods exacerbates gastrointestinal distress. Rather than fermenting safely in the large intestine, fiber ferments prematurely in the small intestine, producing intense gas, severe bloating, micro-perforations (leaky gut syndrome), and systemic symptoms.



1. The Antibiotic Cycle: SIBO sufferers treated with broad-spectrum antibiotics often experience temporary relief due to non-selective bacterial eradication. However, antibiotics decimate both pathogenic and beneficial flora while leaving fungal species untouched. This enables opportunistic yeasts, particularly Candida, to overgrow. Deprived of microbial diversity, the patient suffers recurrent infections, antimicrobial resistance, heightened food allergies, and autoimmune conditions.

The Gut-Brain Axis and Neurological Pathologies

Physiology of the Enteric Nervous System

The gastrointestinal tract operates as a primary neuroendocrine organ, often termed the “second brain.” During embryonic development, tissue folds separate into the Central Nervous System (CNS) and the Enteric Nervous System (ENS), remaining structurally connected via the vagus nerve (the tenth cranial nerve).

SUBSCRIBE

www.naturefresh.co.za

Fresh Talk

FRESH

with NATURE FRESH

Natural Solutions to Health Questions



The vast majority of the body's total serotonin—the critical neurotransmitter regulating mood, impulse control, sleep, and aggression—is synthesized and housed within the gut mucosa, not the brain. Consequently, localized gut inflammation, dysbiosis, or lectin-induced tissue erosion alters central neurochemistry directly via vagal signaling.

Early Microbiome Development

Exposure to normal microflora during critical developmental windows in infancy permanently programs gene expression pathways governing learning, memory, motor function, and stress resilience.

* Introducing complex grain products, infant cereals, or gluten prior to complete digestive tract maturation induces mucosal damage, allergenic sensitization, and microflora depletion.

* Infantile microbial disruption establishes persistent neurodevelopmental and behavioral alterations that cannot be fully corrected by reintroducing beneficial bacteria in adulthood.

Psychiatric and Neurological Conditions

Gluten ingestion and WGA bioaccumulation are linked to severe psychiatric and motor deficits:

NEUROLOGICAL & PSYCHIATRIC MANIFESTATIONS	
Condition	Pathophysiological Connection
Gluten Ataxia	Autoimmune cerebellar damage causing gait disturbance, loss of motor coordination, and vertigo. (40% of idiopathic ataxia cases stem directly from gluten).
Schizophrenia	Exorphin/gliadorphin peptides cross blood-brain barrier; historical epidemiological studies link grain diets directly to schizophrenia prevalence.
Depression & Anxiety	Inhibition of intestinal serotonin synthesis, circulating anti-gliadin antibodies, and vagal inflammatory signaling.
Cognitive Impairment	WGA-mediated myelin degradation, central excitotoxicity, and brain fog.

Historically, isolated populations consuming traditional non-grain diets (such as indigenous groups in Papua New Guinea, the Solomon Islands, and Micronesia) demonstrated near-zero chronic schizophrenia cases (2 individuals out of >65,000 examined). Following the introduction of Western wheat, barley beer, and processed grains, schizophrenia incidence rose to match Western levels.

OFFICIAL DISCLAIMER: All material provided is only information and may not be construed as medical advice. Only a doctor or an appropriate health practitioner is qualified to make a diagnosis and provide the necessary treatment and medications. Products discussed are not intended to diagnose, cure or prevent any form of disease.

SUBSCRIBE

www.naturefresh.co.za

Fresh Talk

FRESH

with NATURE FRESH

Natural Solutions to Health Questions

Genetic and Blood Type Specificity

The Peter D'Adamo Blood Type Framework

Susceptibility to gluten and dietary lectins is governed by inherited genetic markers, specifically blood type antigens and secretor status. Lectins in mismatched foods react directly with blood type surface antigens, provoking cellular agglutination, digestive dysfunction, and autoimmune cascades.

BLOOD TYPE LECTIN & GRAIN INCOMPATIBILITY		
Blood Type Group	Wheat Tolerance	Dietary Incompatibilities & Guidelines
Type A Secretor	Tolerated (A1 subset)	Possesses friendly digestive antigens for wheat; must still avoid raw bran.
Type A Non-Secretor (A2)	Intolerant	Highly prone to rapid, morbid obesity and metabolic failure from wheat.
Type O Secretor	Highly Intolerant	Prone to severe anaemia, joint pain, bowel sluggishness. Chickpeas tolerated but not lentils or corn.
Type O & B Non-Secretors	Intolerant	Severe reactions to wheat gluten; soy protein acts as a damaging lectin.

Crucially, Blood Type A Secretors (A1) represent the minority of the global population capable of properly digesting wheat without immediate antigenic conflict. Because Type A1 individuals often tolerate wheat, they frequently introduce standard wheat products to family members of differing blood types, causing undiagnosed chronic illness in non-secretors or Type O/B relatives.

Allergenic Addiction Mechanism

When individuals consume foods containing lectins or gluten proteins incompatible with their blood type, the digestive system experiences micro-trauma. To buffer this toxic stress, the body releases endogenous opiate peptides (opioids).

This temporary narcotic effect numbs initial digestive distress, creating a psychological and physiological addiction to the offending food (e.g., intense cravings for bread, pastries, sugar, and fat). Over time, this cycles into fat cell swelling, fluid retention, metabolic suppression, and chronic fatigue.

OFFICIAL DISCLAIMER: All material provided is only information and may not be construed as medical advice. Only a doctor or an appropriate health practitioner is qualified to make a diagnosis and provide the necessary treatment and medications. Products discussed are not intended to diagnose, cure or prevent any form of disease.

SUBSCRIBE

www.naturefresh.co.za

Fresh Talk

FRESH Talk

Natural Solutions to Health Questions

with NATURE FRESH

Comprehensive Diagnostic and Identification Protocols

Diagnostic Testing Options

Identifying gluten intolerance and celiac disease requires precise diagnostic pathways:

* Gastroenterological Panel & Small Bowel Biopsy: Essential for diagnosing full celiac disease. Biopsies inspect the small intestine for flattened, damaged villi. Note: Patients must be actively consuming gluten prior to blood testing or intestinal biopsy to produce valid antibody/histological markers.

* ImuPro 100 Delayed Food Allergy Test (IgG Type III): A comprehensive delayed food allergy panel analyzing 100 IgG antibody reactions. Utilized to pinpoint specific non-celiac grain and food sensitivities without medical prescription delays.

* Bio-Feedback Testing: Alternative functional evaluation utilizing devices such as SCIO to screen for subtle bio-energetic sensitivities.

Hidden Dietary and Non-Dietary Sources

Gluten contamination extends far beyond standard bakery items (bread, pasta, cereal, crackers, biscuits, waffles, pancakes, muffins).

HIDDEN SOURCES OF GLUTEN	
Category	Products / Ingredients
Hidden Culinary & Processed Foods	Soy sauce, beer, commercial broths, gravies, coffee creamers, flavored teas, fish sticks, imitation seafood, processed hot dogs, meat substitutes, seasoned chips, candy.
Grains Containing Gluten	Wheat, spelt, rye, barley, kamut, bulgur, couscous, cross-contaminated oats.
Non-Dietary / Consumer Goods	Chewing gum, toothpaste, cosmetic lotions.
Safe Exception	Pure Wheatgrass Juice Powder (harvested from jointless baby grass <6 inches tall, prior to gluten protein formation in the grain head).

OFFICIAL DISCLAIMER: All material provided is only information and may not be construed as medical advice. Only a doctor or an appropriate health practitioner is qualified to make a diagnosis and provide the necessary treatment and medications. Products discussed are not intended to diagnose, cure or prevent any form of disease.

SUBSCRIBE

www.naturefresh.co.za

Fresh Talk

FRESH Talk

Natural Solutions to Health Questions

with NATURE FRESH

Complete A-Z Symptom and Pathology Checklist

Gluten toxicity manifests across a vast spectrum of acute and chronic medical conditions:

- * Gastrointestinal: Dysfunctional bowel, Irritable Bowel Syndrome (IBS), Crohn's disease, ulcerative colitis, chronic abdominal pain, severe gas, bloating, constipation, alternating diarrhoea, gastroesophageal reflux, recurrent mouth sores/stomatitis, pancreatic disorders, gallbladder disease.
- * Metabolic & Endocrine: Rapid unexplainable weight loss or stubborn weight gain, enlarged abdominal girth, insulin resistance, Type 1 diabetes, Addison's disease, hyperthyroidism, hypothyroidism (Hashimoto's), hormonal imbalances, amenorrhea, delayed menarche.
- * Neurological & Psychiatric: Headaches, severe migraines, brain fog, inability to concentrate, memory impairment, depression, anxiety, mood swings, bipolar disorder, schizophrenia, seizures, epilepsy, ataxia, peripheral neuropathy, tingling numbness in extremities.
- * Musculoskeletal: Systemic joint pain, morning stiffness, rheumatoid arthritis, muscle aches, eroded joints, loss of glucosamine, osteoporosis, osteopenia, osteomalacia, pathologic fractures, dental enamel hypoplasia.
- * Dermatological: Dermatitis herpetiformis, severe eczema, psoriasis, alopecia (hair loss), vitiligo, unexplained skin eruptions, poor skin tone.
- * Immunological & Autoimmune: Chronic fatigue syndrome, systemic lupus erythematosus, Multiple Sclerosis (MS), Sjögren syndrome, Non-Hodgkin's lymphoma, autoimmune hepatitis, primary biliary cirrhosis, Total IgA deficiency.
- * Reproductive & Urological: Infertility, recurrent spontaneous abortions/miscarriages, low birth-weight infants, weak bladder, nocturnal enuresis (bedwetting).
- * Systemic / Vascular: Persistent iron-deficiency anaemia, shortness of breath, asthma, elevated inflammatory markers, water-logging/edema.

OFFICIAL DISCLAIMER: All material provided is only information and may not be construed as medical advice. Only a doctor or an appropriate health practitioner is qualified to make a diagnosis and provide the necessary treatment and medications. Products discussed are not intended to diagnose, cure or prevent any form of disease.

SUBSCRIBE

www.naturefresh.co.za

Fresh Talk

FRESH


Natural Solutions to Health Questions

with **NATURE FRESH**



Strategic Dietary Protocols and Solutions

To resolve gluten-mediated disease, individuals must transition away from ultra-processed commercial products toward whole-food nourishment.



The 1-Month Gluten-Free Challenge

A 30-day absolute elimination protocol serves as both a diagnostic tool and a therapeutic baseline.

- * Zero-Tolerance Policy: Complete removal of wheat, rye, barley, and oats. Ingesting as little as 6 grams of gluten (equivalent to a fraction of a slice of bread) can trigger systemic autoimmune activity lasting weeks.
- * Whole-Food Substitutions: Focus on naturally gluten-free whole foods, such as meats, poultry, wild seafood, eggs, non-starchy vegetables, seeds, and healthy natural fats.
- * Baking Alternatives: Where baking is required, utilize single-ingredient, blood-type-compatible whole flours (e.g., sweet potato flour, yellow pea flour, chickpea flour, or butter bean blends bound naturally with fresh eggs) rather than commercially packaged starch-heavy mixes.

OFFICIAL DISCLAIMER: All material provided is only information and may not be construed as medical advice. Only a doctor or an appropriate health practitioner is qualified to make a diagnosis and provide the necessary treatment and medications. Products discussed are not intended to diagnose, cure or prevent any form of disease.

SUBSCRIBE

www.naturefresh.co.za

Fresh Talk

FRESH

Natural Solutions to Health Questions

with NATURE FRESH



Specialized Dietary Interventions

1. Grain-Free Healthy Keto: Eliminates all grains (corn, rice, wheat, oats, quinoa) alongside refined tubers (potatoes, tapioca) to suppress hyper-insulinaemia, reverse fatty liver, halt advanced glycation, and eliminate systemic gluten exposure.
2. The Carnivore Protocol: Recommended for patients suffering from severe SIBO, micro-perforations, extreme bloating, or total inability to ferment vegetable fiber. Consisting entirely of meat, animal fats, and eggs, it removes all plant lectins, WGA, and fermentable fibers, giving the small intestinal mucosa time to repair.
3. Blood Sugar Stabilization (MEGASLIM Strategy): To prevent cortisol spikes, adrenal exhaustion, and junk food cravings during recovery, patients should consume small, protein-rich meals or snacks every 3–4 hours. This practice supports glycogen release from the liver without triggering insulin surges.

Microbiome Restoration

Correcting historical gut damage requires actively repopulating lost microbial diversity:

- * Fermented Vegetables: Daily consumption of raw unpasteurized sauerkraut, kimchi, pickled vegetables, and lacto-fermented pickles.
- * Fermented Dairy (if tolerated): Consumption of raw cheeses, traditional kefir, and unsweetened plain yogurt to re-introduce beneficial bacterial strains.
- * Targeted Probiotics: High-potency broad-spectrum multi-strain probiotics (e.g., PROBIOTIC-15 formulations) to re-establish normal microbial balance, displace pathogenic yeast/candida, and restore mucosal integrity.

OFFICIAL DISCLAIMER: All material provided is only information and may not be construed as medical advice. Only a doctor or an appropriate health practitioner is qualified to make a diagnosis and provide the necessary treatment and medications. Products discussed are not intended to diagnose, cure or prevent any form of disease.

SUBSCRIBE

www.naturefresh.co.za

Fresh Talk

FRESH Talk

with

NATURE
FRESH

Natural Solutions to
Health Questions



OFFICIAL MEDICAL DISCLAIMER

BEFORE YOU BEGIN: A WORD OF CAUTION: This information is not intended as medical diagnosis or treatment. Please consult an appropriate health practitioner for advice if you have a medical problem, as we are not in a position to do so. We merely provide nutritional guidelines that you can learn to establish a suitable eating plan for good results.

It is your responsibility to let your doctor know if you wish to adjust your current medications. Please check up on any potential drug reactions and intolerance you may have to any of the herbs and spices and fresh fruit and vegetables. It is best to consult a health professional that can help you monitor your progress, should you need allergy testing, your blood group identified or advice on supplementation, herbal & homeopathic remedies or medications.

Official DISCLAIMER: *This unregistered medicine has not been evaluated by the SAHPRA for its efficacy, quality, safety or intended use.*

Source References:

<https://www.naturefresh.co.za/Article/223>

<https://www.youtube.com/watch?v=u8YFHAAuC7A>

<https://www.naturefresh.co.za/Article/232>

<https://www.youtube.com/watch?v=XiMxWUF6k04>

<https://www.naturefresh.co.za/Article/247>

<https://www.youtube.com/watch?v=PJ2cW7mF9RE>

<https://www.naturefresh.co.za/Article/249>

<https://www.naturefresh.co.za/Page/Disclaimer>

<https://www.naturefresh.co.za/Article/373>

OFFICIAL DISCLAIMER: All material provided is only information and may not be construed as medical advice. Only a doctor or an appropriate health practitioner is qualified to make a diagnosis and provide the necessary treatment and medications. Products discussed are not intended to diagnose, cure or prevent any form of disease.